# REPORT ON THE SURVEY OF NORTHERN LAKE NYASA

# REPORT ON THE SURVEY OF Northern Lake Nyasa 1954-55

by the

Joint Fisheries Research Organization

E

# P. B. N. JACKSON, M.Sc.

Joint Fisheries Research Organization, Samfya, Northern Rhodesia

T. D. ILES, B.Sc.

Fisheries Laboratory, Ministry of Agriculture and Fisheries, Lowestoft

D. HARDING, B.Sc.

Joint Fisheries Research Organization, Samfya, Northern Rhodesia

AND

G. FRYOR, Ph.D. 🗐

Freshwater Biological Association, Ambleside

1963 PRINTED AND PUBLISHED BY THE GOVERNMENT PRINTER ZOMBA, NYASALAND

#### CONTENTS

| Foreword | •• | •• | •• | •• | <br>•• | •• | 1 |
|----------|----|----|----|----|--------|----|---|
|          |    | _  |    |    |        |    |   |

#### CHAPTER ONE

#### INTRODUCTORY

|    | <b>INTRODUCTION</b> | ••      | ••        | ••      | ••    | ••  | ••  | J |
|----|---------------------|---------|-----------|---------|-------|-----|-----|---|
| 2. | SUMMARY OF RECOI    | MMENDAT | TIONS AND | CONCLUS | SIONS | • • | • • | 5 |

#### CHAPTER TWO

### STUDIES ON THE HYDROLOGY OF LAKE NYASA AND ASSOCIATED RIVERS

|    | DESCRIPTION OF THE LAKE                 |      |    |    | 10 |
|----|---|------|----|----|----|
| 2. | METEOROLOGICAL DATA AFFECTING THE LAKE  | ••   | •• | •• | 11 |
| 3. | Analyses of Inflows into the Lake       |      |    | •• | 12 |
| 4. | WATER TEMPERATURES AND CHEMISTRY OF THE | Lake | •• | •• | 14 |

#### CHAPTER THREE

#### STUDIES ON THE INVERTEBRATES, WITH ESPECIAL REFERENCE TO CRUSTACEA

| 1. | Fri | BE-LIVING CRUSTACEA       | ••         | ••       | ••      | ••       |      | 45 |
|----|-----|---------------------------|------------|----------|---------|----------|------|----|
| 2. | PAI | RASITIC CRUSTACEA:        |            |          |         |          |      |    |
|    | (a) | Quint III                 | •••        | ••       | ••      | •••      | ••   | 50 |
|    | (b) | Key to the species of par | asitic Cop | epoda an | d Branc | hiura of | Lake | 50 |
|    | • • | Nyasa                     | ••         | ••       | ••      | ••       | ••   | 52 |
| 3. | Nơ  | TES ON SOME OTHER INV     | ERTEBRAT   | e Fauna  | • •     | ••       | ••   | 53 |

#### CHAPTER FOUR

# THE FISHES OF NYASALAND, WITH ESPECIAL REFERENCE TO NORTHERN LAKE NYASA

| 1. | Stu | DIES ON SYSTEMATICS AND     | Ecolo  | GY:       |                |    |     |     |
|----|-----|-----------------------------|--------|-----------|----------------|----|-----|-----|
|    | A.  | Introduction                |        | ••        | ••             | •• | • • | 55  |
|    | B.  | Broad Ecological Zones in   | the Ny | asan Area | ••             |    | • • | 55  |
|    |     | Check-list of the Fishes of |        |           | ••             | •• | ••  | 58  |
|    |     | Notes on Zambezi Fishes     | ••     | ••        | ••             |    | ••  | 92  |
| 2. |     | res on the Biology of H     | BAGRUS | MERIDION  | AL <b>I</b> S: |    |     |     |
|    | A.  | Breeding                    | ••     | ••        | ••             | •• |     | 95  |
|    |     | (1) Sex Ratio               | ••     | ••        | ••             | •• | ••  | 95  |
|    |     | (2) Breeding Cycle          | • •    | ••        |                |    | • • | 96  |
|    |     | (3) Breeding Behaviour      | ••     | ••        |                | •• |     | 99  |
|    | B.  | Feeding                     | ••     | ••        | ••             | •• |     | 100 |

9

# 3. THE GENERAL BIOLOGY OF UTAKA:

| Systematics                |  |   |   |   |   | 102   |
|----------------------------|--|---|---|---|---|---|
| Habitats and Abundance .   | •  | ••  | ••  | ••  | ••  | 102   |
| Feeding                    | •  | ••  | ••  | ••  | ••  | 106   |
| Breeding and Growth Rate   |  |   | ••  | ••  | ••  | 107   |
| (I) Haplochromis quadri    | maculatu   | s   | ••  |   |   | 107   |
| (II) Haplochromis virgind  | alis   | ••  |   | • •   |   | 109   |
| (III) Haplochromis pleuros | stigmoi <mark>d</mark> e   | 5   | • •   |   |   | 110   |
| (IV) Haplochromis mloto    |  | ••  | ••  | • •   | ••  | 111   |
| (V) Haplochromis barleyi   |  | ••  | ••  | ••  |   | 111   |
| (VI) Other Species .       | •  | ••  | ••  | ••  |   | 112   |
|                            | Habitats and Abundance<br>Feeding<br>Breeding and Growth Rate<br>(I) Haplochromis quadri<br>(II) Haplochromis virgind<br>(III) Haplochromis pleuros<br>(IV) Haplochromis mloto<br>(V) Haplochromis barleyi | <ul> <li>Habitats and Abundance</li> <li>Feeding</li> <li>Breeding and Growth Rate <ul> <li>(I) Haplochromis quadrimaculatu</li> <li>(II) Haplochromis virginalis</li> <li>(III) Haplochromis pleurostigmoide</li> <li>(IV) Haplochromis mloto</li> <li>(V) Haplochromis barleyi</li> </ul> </li> </ul> | Habitats and Abundance          Feeding          Breeding and Growth Rate          (I)       Haplochromis quadrimaculatus         (II)       Haplochromis virginalis         (III)       Haplochromis pleurostigmoides         (IV)       Haplochromis mloto         (V)       Haplochromis barleyi | Habitats and Abundance          Feeding          Breeding and Growth Rate          (I) Haplochromis quadrimaculatus          (II) Haplochromis virginalis          (III) Haplochromis pleurostigmoides          (IV) Haplochromis mloto          (V) Haplochromis barleyi          (VI) Other Species | Habitats and Abundance          Feeding          Breeding and Growth Rate          (I) Haplochromis quadrimaculatus          (II) Haplochromis virginalis          (III) Haplochromis pleurostigmoides          (IV) Haplochromis mloto          (V) Haplochromis barleyi          (VI) Other Species | Habitats and Abundance          Feeding          Breeding and Growth Rate          (I) Haplochromis quadrimaculatus          (II) Haplochromis virginalis          (III) Haplochromis pleurostigmoides          (IV) Haplochromis mloto          (V) Haplochromis barleyi          (VI) Other Species |

### CHAPTER FIVE

# THE FISHERIES OF THE NORTHERN LAKE

| 1. | Тне | Gill-Net    | FISHERY | OF | THE | Northern | LAKE: |  |
|----|-----|-------------|---------|----|-----|----------|-------|--|
|    | Α   | Introductio | n       |    |     |          |       |  |

| А. | Introduction          | ••                  | ••          | ••       | • •       | ••  |    | 118         |
|----|-----------------------|---------------------|-------------|----------|-----------|-----|----|-------------|
| В. | Procedure             | • •                 | ••          | ••       | ••        | ••  | •• | 120         |
|    | I. Descri             | ption of a          | Gill-Net    | ••       |           | ••  |    | 120         |
|    | II. Some              | <b>Fheor</b> etical | Considera   | ations   |           |     | •• | 120         |
|    | III. Mount            | ing                 |             |          |           | ••  | •• | 121         |
|    | IV. Suitab            | le Lengths          | for Mount   | ting     | ••        |     |    | 122         |
|    | V. Optim              |                     |             | ••       |           | ••  |    | 125         |
|    | VI. Optimi            | -                   |             | ••       |           |     |    | 128         |
|    |                       | epth Distri         | -           |          |           |     | •• | 128         |
|    |                       | esults              |             |          |           |     | •• | 120         |
|    | VII. Laying           | of Nets             | ••          | ••       |           |     | •• | 129         |
|    | VIII. Care a          |                     | nance of G  |          |           | ••  | •• | 130         |
| C. | The Experime          |                     |             |          |           | ••  | •• | 130         |
|    | I. General            |                     |             | ,        | ••        | ••  | •• | 131         |
|    | II. Sm <b>a</b> ll-me | sh Nets             |             | ••       | ••        | ••  | •• | 131         |
|    | (a) $2\frac{1}{2}$ in |                     |             |          | ••        | ••  | •• | 132         |
|    | • • -                 | h Nets              |             | ••       | ••        | ••  | •• |             |
|    | III. Large-me         |                     | ••          | ••       | ••        | ••  | •• | 135         |
|    |                       | eral                | ••          | ••       | ••        | ••  | •• | 136         |
|    | .,                    |                     |             |          | ••        | ••  | •• | 136         |
|    |                       | acteristics         |             |          | ll-Nets   | ••  | •• | 138         |
|    | (i)                   |                     | Weight of   |          | ••        | ••  | •• | 138         |
|    | (ii)                  | Relative            | Occurrence  | e of Spe | cies      | ••  | •• | <b>13</b> 8 |
|    | (iii)                 | Variation           | in Catch    | through  | out the Y | ear | •• | 139         |
|    | (iv)                  | Length of           | f Fishing U | Jnit     | ••        | ••  | •• | 140         |
|    | (v)                   | Depth of            | Net         | ••       | ••        | ••  | •• | 141         |
|    | (vi)                  | Twine ty            | pe and thi  | ckness   | ••        | ••  | •• | 142         |
|    | (vii)                 | Mesh size           | ••          | ••       | ••        | ••  | •• | 144         |
|    |                       |                     |             |          |           |     |    |             |

| 2.         | Тне | UTAKA FISHERY IN THE NORTHE          | RN LAP  | KE :       |           |     |     |
|------------|-----|--------------------------------------|---------|------------|-----------|-----|-----|
|            | A.  | Introduction                         | ••      | ••         | ••        | ••  | 145 |
|            | В.  | The present Fishery in the Norther   | rn Lake | •••        | ••        | ••  | 146 |
|            |     | I. Methods                           | ••      | ••         | ••        | ••  | 146 |
|            |     | (a) The Chilimila Net                | ••      | ••         | ••        | ••  | 146 |
|            |     | (b) The Kawelekete Net               | ••      | ••         | ••        | ••  | 150 |
|            |     | (c) The Ngongongo Net                | ••      | ••         | ••        | ••  | 151 |
|            |     | (d) Beach Seines                     | ••      | ••         | ••        | ••  | 151 |
|            | C.  | Experimental Fishing                 | ••      | ••         | ••        | ••  | 152 |
|            |     | I. Ring Netting                      | ••      | ••         | ••        | ••  | 152 |
|            |     | II. Floating Trawl                   | ••      | ••         | ••        | ••  | 153 |
|            |     | III. High-Speed Self-opening Traw    | vl      | ••         | ••        | ••  | 153 |
|            |     | IV. Small-mesh Chilimila             | • •     | ••         | ••        | ••  | 153 |
|            |     | (a) Catch Data                       | ••      | ••         | ••        | ••  | 153 |
| 3.         | DAM | ABO FISHING                          | ••      | ••         | ••        | ••  | 158 |
| <b>4</b> . | Rıv | ERS AND THE BARILIUS FISHERY:        |         |            |           |     |     |
|            | A.  | General                              | ••      | ••         |           | ••  | 159 |
|            | В.  | The Spawning Migration and Afric     | can Riv | er Fishery | ••        | ••  | 160 |
|            | C.  | African Fisheries for Barilius durin |         |            | Migration | ••  | 161 |
|            |     | I. The Spawning Fishery in the       | Luweya  | a River    | ••        | ••  | 161 |
|            |     | II. Other Barilius Fisheries         | ••      |            | ••        | ••  | 165 |
|            | D.  | Discussion and Recommendations       |         | ••         | ••        | ••  | 165 |
| 5.         | Тня | E FISHERY FOR USIPA:                 |         |            |           |     |     |
|            | А.  | Introduction                         | ••      | ••         | • •       | ••  | 167 |
|            | В.  | The Chiwu Method                     | ••      | ••         | ••        | ••  | 168 |
|            | С.  | The Beach Seine                      | ••      | ••         | ••        | ••  | 169 |
|            | D.  | The Usikite Net                      | • •     | ••         | ••        | ••  | 171 |
|            | E.  | Discussion                           | ••      | ••         | ••        | • • | 171 |
|            |     |                                      |         |            |           |     |     |

#### MAPS:

1 to 6 as folders in the back of the report.

#### LIST OF ILLUSTRATIONS

- Plate I A. The laboratory, Nkata Bay.
  - B. The research launch moored at Nkata Bay.
  - C. Method of mounting gill-nets to head rope.

D. Floats crushed by pressure in the deep gill-net fishery. The three on the left are unused.

#### Plate II

#### MAJOR ECOLOGICAL ZONES

- A. A sandy sheltered bay in the northern lake; habitat for zone V fishes at Nkata Bay.
- B. A small sandy beach on the open northern lake, sharply adjacent to the more usual rocky shore; habitats for fishes of zones V and VI respectively near Nkata Bay.
- C. A rocky shore with some intermediate rock/sand area in the foreground; habitats for fishes of zones VI and VII respectively near Nkata Bay.
- D. Rock/sand zone; habitat for zone VII fishes near Nkata Bay.
- E. Dhows and dug-outs at Fort Maguire; the weedy stagnant gutway in the foreground provides a typical habitat for zone III fishes in the southern lake, while sheltered zone IV conditions are beyond.
- F. A sheltered bay in the southern lake; zones V and VI at Monkey Bay.
- G. Sheltered zone V conditions in the northern lake; the anchorage at Bana.
- H. A slow-flowing reedy river in its lower reaches, habitat for zone III fishes; the Limpasa River ten miles inland from Nkata Bay.

Plate III

#### THE DEEP GILL-NET FISHERY

- A. A large haul of Tilapia and Clariids; south-west arm.
- B. Setting a deep gill-net, Nkata Bay.
- C. Gill-nets and fish after hauling from 60-70 metres, Nkata Bay.
- D. Bathyclarias nyasensis, B. longibarbis and Bagrus meridionalis in gillnets hauled from 60-70 metres, Nkata Bay. Note the distended abdomen of the Bagrus caused by the rapid change in pressure.
- Plate IV Four species of Utaka from a chirimila at Nkata Bay. From top to bottom:
  - A. Haplochromis H, H, H,
  - B. Bathyclarias gigas, Nkata Bay.
  - C. Bathyclarias filicibarbis, Nkata Bay.
  - D. Haplochromis heterotaenia Nkata Bay.

#### AFRICAN FISHERIES

- Plate V
- A. The gorge at Chiwandama, Luweya river, where Sanjika (Barilius microcephalus) are caught on their upstream breeding migration.
  - B. Breeding Sanjika being lifted from slack water in crevice of the rock, where they rest before attempting to move further up the rapids. Note the construction of the "Khombe" net.
  - C. "Mono" basket set in deep water to catch Synodontis njassae. Usually baited with dough as fish bait attracts crabs. Near Nkata Bay.
  - D. Arrangement of canoe for Usipa fishing. The torch-bearer is forward with the netsman close to him, while the paddler sits astern.

#### Plate VI

- A, B, C. A seasonal fishery for breeding *Rhamphochromis* (probably *R. longiceps*) at Bana near the entrance to the lagoon (obstructed by weir, C). Small seines are hauled by the two men.
- D. A large beach seine being hauled in the south-west arm.
- E, F. Fishing in Kambwe lagoon near Karonga with scoop nets, and baskets are stabbed vertically downwards, trapped fish being removed through an aperture halfway up.
- G. A "Mono" basket trap filled with crabs, the result of baiting with fish bait, Nkata Bay.

## LIST OF FIGURES

# Chapter Two

|      |     |  | facing<br>page                          |
|------|-----|--|---|
| Fig. | 1.  | Lake Nyasa after E. L. Rhoades and W. B. Phillips 1901 (from van<br>Meel 1954). The shaded area on the chart indicates the part<br>of the lake which lies below sea level  | 10                                      |
| Fig. | 2.  | Rainfall at Nkata Bay between November, 1953, and August, 1955   | 12                                      |
| Fig. | 3.  | Profiles of oxygen, pH and temperature in the South Bay, Nkata Bay, March, 1954  | 14                                      |
| Fig. | 4.  | Station I: Nkata Bay 28.vii.54. Distribution of temperature, oxygen silicate and phosphate in the top three hundred metres at this station   | 22 et<br>seq. to<br>p. 23               |
| Fig. | 5.  | Station I: Nkata Bay 2.ix.54. Temperature and oxygen profiles  | ,,                                      |
| Fig. | 6.  | Temperature distribution at a Station approximately 20 miles NNW. of Likoma Island. 14.ix.54   | "                                       |
| Fig. | 7.  | Thermoclines recorded (a) with a Freidinger reversing thermometer<br>and (b) a thermistor. Readings taken simultaneously at Nkata Bay<br>near Station I, 3.xii.54  | ,,                                      |
| Fig. | 8.  | Distribution of temperature, oxygen and silicate at Station I, Nkata<br>Bay. 25.iii.55   | رر                                      |
| Fig. | 9.  | Temperature profiles on several dates during 1954 and 1955 at Station I, Nkata Bay   | ,,                                      |
| Fig. | 10. | Isotherms at Station I, Nkata Bay, between the months of September, 1954, and October, 1955. Wind speed and barometric pressure are included on the graph for the same period  | ,,                                      |
| Fig. | 11. | Temperature change at seven depths, recorded at Station I, Nkata<br>Bay, between March, 1954, and October, 1955  | دو                                      |
| Fig. | 12. | Annual fluctuation in the phosphate content of water taken from five different levels at the Nkata Bay deep station during 1954 and 1955. Phosphate measured as mg./L. P.  | ,,                                      |
| Fig. | 13. | Temperature profile and silicate distribution at Station I, Nkata Bay, on 7.ii.55  |   |
| Fig. | 14. | Conductivity change (K corr) at 0, 50, 100, 200, and 300 metres, at Station I, near Nkata Bay  |   |
| Fig. | 15. | Plankton volumes stained from the top hundred metres using coarse (60 m.p.i.) and fine (120 m.p.i.) nets. Volumes given are those sedimented from 7,070 litres of lake water. Silicate mean values in the top fifty metres are superimposed on the histogram. Station I, Nkata Bay, March, 1954, to October, 1955  | ""                                      |
| Fig. | 16. | Usisya Deep Station (over 385 fathoms). Profiles of temperature, oxygen, silicate and phosphate down to 600 metres. 21.vii.55  | ,,                                      |
| Fig. | 17. | Bridge diagram and calibration curve of a thermistor set used at Nkata Bay. Lettering: $T = $ Stantel Type Thermistor (2311/300); $S = $ Helical Potentiometer of 1,000 ohms resistance; Resistances: $R_1 = 1,600$ ohms; $R = 2,500$ ohms; $G = $ galvonometer 3,000 ohms resistance with maximum sensitivity at the centre zero. The battery used in the circuit was a 1.34 v. Ruben Mallory Battery RMB-3 $\dots$ | .,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, |

#### **Chapter Three**

facing page

The Planktonic Crustacea of Lake Nyasa Fig. 1. 1. Daphnia lumholtzi. 2. Bosmina longirostris. 3. Bosminopsis dietersi. 4. Diaphanosoma excisum. 5. Diaptomus (Thermodiaptomus) mixtus. 6. D. mixtus (another view). 7. Diaptomus (Tropodiaptomus) kraepelini. 8. Mesocyclops neglectus. 9. Mesocyclops leukarti. (In part original, in part modified from Sars, Birge and Gunther.)

Examples of the Crustacean parasites of the fishes of Lake Nyasa ... Fig. 2. 1. Chonopeltis inermis. 2. Argulus africanus. 3. Afrolernea longicollis. 4. Lamproglena nyasae. 5. Ergasilus cunningtoni. 6. Lernea bagri. (After Harding and others original or simplified from Fryer.)

#### **Chapter Four**

#### Section 2

| Fig. | 1. | Bagrus meridionalis. Variation of percentage males with length | ر 100 et                |
|------|----|--|-------------------------|
| Fig. | 2. | B. meridionalis. Breeding cycle of female                      | l con to                |
| Fig. | 3. | B. meridionalis. Percentage males caught throughout the year   | D. 101                  |
| Fig. | 4. | B. meridionalis. Stomach contents                              | <b>J P</b> . <b>1</b> . |

#### Section 3

| Fig. | 1. | Haplochromis quadrimaculatus. Length frequency. Breeding fish | ך<br>ר    |
|------|----|---|-----------|
| Fig. | 2. | H. quadrimaculatus. Length frequency diagrams                 |           |
| Fig. | 3. | H. quadrimaculatus. Growth of 1954 year class                 | 112 et    |
| Fig. | 4. | H. virginalis. Length frequency breeding fish                 | ≻ seq. to |
| Fig. | 5. | H. virginalis. Length frequency diagrams (Two pages)          | p. 113    |
| Fig. | 6. | H. virginalis. Growth of 1954 year class                      |           |
| Fig. | 7. | H. borleyi. Length frequency of breeding fish                 | J         |

#### **Chapter Five**

#### Section 1

| Fig. 1.   | Diagram of stretched length of netting   |  |  |  |  |  |  |  |  |
|-----------|--|--|--|--|--|--|--|--|--|
| Fig. 2.   | Correct method of mounting a gill-net 122 et                                     |  |  |  |  |  |  |  |  |
| Fig. 3.   | Single mesh of net at its maximum fishing area > seq. to                         |  |  |  |  |  |  |  |  |
| Fig. 4.   | Relation between area of gill-net and method of mounting p. 123                  |  |  |  |  |  |  |  |  |
| Fig. 5.   | Method of mounting weights   |  |  |  |  |  |  |  |  |
| Fig. 6.   | Distribution of fish in 5" 26 mesh gill-net                                      |  |  |  |  |  |  |  |  |
| Fig. 7.   | Relation between fish and mesh shape in two types of gill-net fishery $> 128$ et |  |  |  |  |  |  |  |  |
| Fig. 8.   | Determination of depth profile of gill-net seq. to                               |  |  |  |  |  |  |  |  |
| Fig. 9.   | Typical depth profile $\dots$ $\dots$ $\dots$ $p. 129$                           |  |  |  |  |  |  |  |  |
| Fig. 10.  | Variation in Nos. Fish/Unit effort throughout year 140                           |  |  |  |  |  |  |  |  |
| (a & 1    |  |  |  |  |  |  |  |  |  |
| Fig. 11.  | Frequency histogram of catches with nylon and flax nets 142                      |  |  |  |  |  |  |  |  |
| Section 2 |  |  |  |  |  |  |  |  |  |

#### Section 2

| Fig. | 1. | Chilimila net | • • | •• | •• | •• | •• | •• | 146 |
|------|----|---------------|-----|----|----|----|----|----|-----|
|------|----|---------------|-----|----|----|----|----|----|-----|

50

#### Maps and Charts

- Map 1. Contours of Lake Nyasa, land above lake level showing contours and rivers entering the lake, sampling stations.
- Map 2. Chart of Lake Nyasa, compiled by J. G. Pike, including soundings taken during several cruises in the Northern Lake Nyasa and soundings after Rhoades.
- Map 3. Likoma and Chisumulu Islands showing soundings and indicating the location of the main Utaka grounds.
- Map 4. Mbamba Bay, Likoma Island, showing fishing grounds.
- Map 5. Enlarged section of the Utaka fishing grounds-Likoma Island. Dotted areas are submerged Virundu.

Map 6. Virundu at Nkata Bay.

с э

6.8